Milwaukee County COVID-19 Data Summary

Milwaukee County COVID-19 Epidemiology Intel Team

This report was updated on October 22, 2020 and includes data through October 20, 2020. Note that data for recent weeks may be under-reported due to pending test results.

Milwaukee County COVID-19 Summary Statistics

Overall Milwaukee County COVID-19 Summary Statistics March 1 - October 20					
	Milwaukee County	City of Milwaukee	Suburbs		
Total tests performed	514,130	335,444	178,686		
Percent positive of all tests performed	8.1%	8.6%	7.1%		
Number of confirmed cases	36,555	25,807	10,748		
Number of hospitalizations	2,769	1,981	788		
Number of deaths	553	351	202		
Case fatality rate	1.5%	1.4%	1.9%		

Weekly Milwaukee County COVID-19 Summary Statistics October 14 - October 20				
	Milwaukee County	City of Milwaukee	Suburbs	
Total tests performed	20,664	12,776	7,888	
Percent positive of all tests performed	11.1%	11.0%	11.2%	
Number of confirmed cases	2,031	1,334	697	
Number of hospitalizations	152	93	59	
Number of deaths	4	2	2	

Total Cases and New Cases

There are now a total of 36555 cases in Milwaukee County, since the first confirmed case on March 6th, 2020. Over the last week, we observed 2031 new confirmed cases in Milwaukee County, including 1334 new cases in the city of Milwaukee. **Figure 1** shows the daily incidence of new cases (bars) and the average daily incidence within the last 7 days (line), which provides a smoothing effect to enhance visualization, for both the city and the county. To indicate a potential reporting delay, we shade the last four days of data and exclude those days from the trend line.

Over the last week, we have seen a continued increase in confirmed cases, particularly in the City of Milwaukee. The highest daily case count since the beginning of the epidemic occurred on October 12, 2020, with 577 cases in the county overall. The highest daily case count over the entire epidemic in the suburbs occurred on October 12, 2020, with a total of 200 cases confirmed, while the highest case count in the city occurred on October 12, 2020, with a total of 377 cases confirmed.




Figure 1: Milwaukee County daily number of COVID-19 cases

Date of specimen collection

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Total Deaths and New Deaths

There are a total of 553 COVID-19 related deaths in Milwaukee County. Over the last week, we observed 4 deaths, with 2 from the City of Milwaukee. **Figure 2** shows the number of daily COVID-19 related deaths among Milwaukee County and City of Milwaukee residents. The overlaid lines show the average daily deaths within the last 7 days for each jurisdiction. Overall, daily deaths have remained relatively stable since mid-July. Deaths peaked on April 8, 2020. Several smaller peaks in deaths are notable since April 8th, with an overall decrease to a low number of deaths beginning in early June.

13 12 Milwaukee County Suburbs 11 City of Milwaukee 10 9 Daily deaths 8 7 6 5 4 3 2 1 Sepon 49/18/18/18/19/ And De De Le St House And An Ant An St Viatiotiatia ce

Figure 2: Milwaukee County COVID-19 daily deaths

Date of death

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


The COVID-19 Growth Rate

The time it takes for the number of cases to double is called the doubling time. **Figure 3** shows doubling times for Milwaukee County, the surrounding Waukesha, Ozaukee and Walworth (WOW) counties, the M7 (7-county) metropolitan area, and the state of Wisconsin. Dotted lines indicate doubling times of 1, 2 3 and 4 days, which are generally associated with a condition of exponential growth. The current doubling time in Milwaukee County is 111.03 days. The current doubling time for WOW counties is 60.95 days. The current doubling time for the state of Wisconsin is 42.78 days. **Figure 4** shows the trend in doubling times for Milwaukee County and the City of Milwaukee as compared to the state, over the course of the epidemic. As illustrated, the epidemic initially doubled more quickly in Milwaukee County and the city, but has since slowed (improved) more in the city and county than in the state as a whole.

Figure 3: Cumulative cases after 20 confirmed Confirmed cases (log10 scale) MKE County 100000 MKE Metro **WOW Counties** 10000 WI State 1000 1-day doubling 2-day doubling 3-day doubling 100 4-day doubling 80 100 120 140 180 200 220 20 40 60 160 Days since 20 confirmed cases

Data source: Wisconsin Department of Health Services Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Figure 4: Trend in doubling times



Data sources: WI Department of Health Services & WI Electronic Disease Surveillance System Created by the Milwaukee County COVID-19 Epidemiology Intel Team

The COVID-19 Reproductive Number

Another way of examining the growth rate of the infection is to examine the reproductive number (R). This number captures the number of new cases that are the result of an existing case. For example, an R of 2 would indicate that each infected person infects 2 new people. The following plots show the change in R over time for Milwaukee County, **Figure 5**, the City of Milwaukee, **Figure 6a**, and Milwaukee County suburbs, **Figure 6b**. Each plot includes key dates related to physical distancing or focused testing campaigns affecting residents. The R for each date is calculated to represent the R for a 7-day period with the start day of that 7-day period represented on the graph.

After the first minimum R value in Milwaukee County observed (R = 0.84 on April 7, 2020), we observed an increase in R to a high of 1.52 on May 8, 2020 and then a decrease to a low of 0.76 in the county on May 21, 2020. The R increased again to a recent high of 1.41 on September 10, 2020. Patterns in the City of Milwaukee are very similar to those in the county overall. Patterns in the suburbs show more fluctuation. The R values for the most recent week are 0.864 for the county, 0.869 in the city, and 0.854 in the suburbs.


5.0 Aug 1 Oct 8 4th of July holiday First distancing Easter & WI state May 11: Two free test sites open Statewide 4.5 mask order begins order 25% assover weekend May 13: WI Supreme Court ruling Reproductive number, capacity 4.0 May 22: Suburbs partially re-open March 25 April 18-20 July 10 Aug 18-Sept 8 HOC testing Safer Several schools May 25: Memorial Day Shorewood 3.5 May 26: City partially re-opens April 24-28 3.0 Smithfield City of MKF - . Labor Day testina mask order begins June 5 2.5 WI election City restaurants 2.0 1.5 1.0 0.5 0.0 Mayor Julion AUGO¹ ~ Oct.08 octo Nax 184 184 5 410 410 410, 5 POL POL POL "V. munumining Mr. Mr. Mr. 50 egegegegeg

Figure 5: One week reproductive number for Milwaukee County

One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


Figure 6a: One week reproductive number for City of Milwaukee



One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Figure 6b: One week reproductive number for Milwaukee County suburbs



One-week window start date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


Demographic Patterns – Age, Sex, Race and Ethnicity

Confirmed cases

COVID-19 cases vary by demographic characteristics. **Figure 7** shows cumulative case plots including confirmed positive cases with an available specimen collection date, plotted by census block group (CBG) median household income, sex, age, and race/ethnicity groups. Most diagnosed cases fall within the ages of 18-79. Of all confirmed cases, 46% are male and 53% are female. The largest number of cases have been identified among the non-Hispanic White population (N = 12965), followed by the Hispanic population (N = 10474), and the Black/AA population (N = 8557). The lower two quartiles of median household income (0 - 35,033, and 0.096) have a larger number of cases than the higher two quartiles (0.097) to 0.097 to 0.096, and 0.096, and 0.096, with the fewest cases identified among the highest income group.

Over the past week, we have observed increases among individuals in all income groups, most age groups, and among non-Hispanic Whites, Hispanics, and non-Hispanic Black/AAs with similar increases for both sexes. The cumulative number of cases among those ages 25-39 (N = 11138) still exceeds the number among the next highest group, those ages 40-59 (N = 9780). In the last few weeks, we saw the number of cases under age 18 (N = 3969) increase to approach the number diagnosed among those 60-79 (N = 4295).

Figure 7: Cumulative confirmed cases in Milwaukee County



Date of specimen collection


Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.


When examined as population-based rates in **Figure 8**, demographic patterns are also apparent. Early in the epidemic, we saw a clear age gradient in population-based rates, with older populations experiencing greater rates. However, in the last months, we have seen rates among the younger, working age groups (18-24, 25-39, and 40-59) exceed the rate of those aged 80+. The rate among those 60-79 now approaches that of those aged 80+. By race and ethnicity, the rate was highest among Black/AA populations until the beginning of May, when we observed a surge among Hispanics resulting in the Hispanic rate (74.51 per 1,000 people) exceeding that among all other racial and ethnic groups. The rate among AIANs (36.8 per 1,000 people) is now slightly higher than that among Black/AAs (34.36 per 1,000 people), although the total number of cases among AIANs (N = 171) is much lower than the number among Black/AAs (N = 8557). The population-based rate among non-Hispanic Whites is relatively low compared to other racial and ethnic groups. Rates are very similar among males and females.

Figure 8: Population based incidence rates in Milwaukee County




Age	N Cases	Population	Rate per 1,000 residents
< 18	3969	231111	17.17
18-24	6266	150895	41.53
25-39	11138	163246	68.23
40-59	9780	230887	42.36
60-79	4295	142783	30.08
80+	1107	35287	31.37

Specimen collection date



Race/Ethnicity*	N Cases	Population	Rate per 1,000 residents
Black or AA	8557	249011	34.36
White	12965	493723	26.26
Hispanic	10474	140575	74.51
Asian	1223	40443	30.24
AIAN	171	4647	36.80
Multiple Races	149	24224	6.15

Specimen collection date



Gender	N Cases	Population	Rate per 1,000 residents
Male	16995	461670	36.81
Female	19471	492539	39.53

Specimen collection date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

Hospitalizations

A total of 2769 individuals have been hospitalized due to COVID-19 in the county. **Figure 9** shows cumulative hospitalizations based on lab specimen collection date (as admission dates are incomplete). The highest number of hospitalizations continues to be among those ages 60-79 (N = 1024). The highest number of hospitalizations have occurred among the Black/AA community (N = 1099), followed by the Non-Hispanic White community (N = 923) and then the Hispanic community (N = 530). Overall, counts are lower among other racial and ethnic groups. Females now slightly outnumber males, comprising 51.6% of all hospitalized cases. More individuals among lower income than higher income groups have been hospitalized, with a clear income gradient observed. Over the last week, we have seen an increase in hospitalizations among all income groups, those 60-79 and 80+, Non-Hispanic Black/AAs and Whites, and both sexes.

Median household income quartile Age 1400 1400 Q1: Lowest < 18 1200 Q2: Low-medium 1200 18-24 Q3: Medium-high 25-39 1000 Q4: Highest 1000 40-59 Cumulative cases among patients who became hospitalized 60-79 800 800 600 600 400 400 200 200 mpmesses to of to gray in in in in in in the 18 18 1 PG Race/Ethnicity* Gender 1400 1400 Black or AA 1200 White 1200 Female Other/Unknown Hispanic 1000 . Asian 1000 AIAN 800 NHOPI 800 Multiple Races Other 600 600 400 400 200 200

Figure 9: Cumulative hospitalizations in Milwaukee County

Date of specimen collection

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team


*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

When examined as population-based rates and case-based rates in **Figure 10**, hospitalization patterns are also apparent by demographic characteristics. Both population- and case-based hospitalization rates exhibit a clear age group gradient, with older age groups experiencing higher rates. By race and ethnicity, population and case-based hospitalization rates are highest among the Black/AA population and the population-based rate is lowest for non-Hispanic Whites. Rates by gender are very similar, with higher hospitalization rates among males. All rates presented are crude rates and only groups with 10 or more


total hospitalized cases are shown. Over the last week, we have seen a concerning uptick in the hospitalization rate among those 80+, and to a lesser degree, those 60-79.

Figure 10: Population and case based hospitalization rates in Milwaukee County




Age	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
< 18	66	0.29	1.66
18-24	102	0.68	1.63
25-39	371	2.27	3.33
40-59	750	3.25	7.67
60-79	1024	7.17	23.84
80+	456	12.92	41.19

Specimen collection date



Race/Ethnicity*	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
Black or AA	1099	4.41	12.84
White	923	1.87	7.12
Hispanic	530	3.77	5.06
Asian	116	2.87	9.48
AIAN	19	4.09	11.11

Specimen collection date



Gender	N Hospitalized Cases	Rate per 1,000 residents	Rate per 100 cases
Male	1338	2.90	7.87
Female	1429	2.90	7.34

Specimen collection date

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

Deaths

There are now a total of 553 confirmed deaths in Milwaukee County, representing a case fatality rate of 1.5%. We observed 4 new deaths over the past week in the county. Mortality patterns differ by demographic characteristics, as shown in **Figure 11**. The largest number of deaths are recorded among those age 60 or older. The largest number of deaths are recorded for males (N = 282) and for non-Hispanic Whites (N = 258) followed by Black/AA residents (N = 199). By income, there are a larger number of deaths among the two lower income groups as compared to the two higher income groups. Deaths among Hispanics remain relatively low.

300 300 Median household income quartile Age Q1: Lowest < 18 250 250 Q2: Low-medium 18-24 Q3: Medium-high 25-39 200 Q4: Highest 200 40-59 60-79 150 150 100 100 50 50 0 APTO Jun.01 ورازار APTO Augon 'sepion Julos 405,00 ,6 101/10/10 bolloglog milning Cumulative deaths 300 300 Gender Race/Ethnicity* Black or AA 250 250 White Female Other/Unknown Hispanic 200 200 Asian AIAN NHOPI 150 150 Multiple Races Other 100 100 50 50 38580000 (5 0, 1,5 0, to to ve of ve of ve of ve of ve of ve of ve

Figure 11: Cumulative deaths in Milwaukee County

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)
Created by the Milwaukee County COVID-19 Epidemiology Intel Team


*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.


In terms of population- and case-based rates shown in **Figure 12**, there is a clear age category gradient, with higher death rates among older populations. Gender-based rates are very similar. Black/AA populations have the highest population and case-based death rates. All rates presented are crude rates and only groups with 10 or more total deaths are shown.

Date of death


Figure 12: Population and case based death rates in Milwaukee County



Age	N Deaths	Rate per 1,000 residents	Rate per 100 cases
18-39	10	0.03	0.06
40-59	64	0.28	0.65
60-79	245	1.72	5.70
80+	234	6.63	21.14



Race/Ethnicity*	N Deaths	Rate per 1,000 residents	Rate per 100 cases
Black or AA	199	0.80	2.33
White	258	0.52	1.99
Hispanic	69	0.49	0.66
Asian	17	0.42	1.39



Gender	N Deaths	Rate per 1,000 residents	Rate per 100 cases
Male	282	0.61	1.66
Female	270	0.55	1.39

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS)

Created by the Milwaukee County COVID-19 Epidemiology Intel Team

*Race and ethnicity were combined into one variable where the Hispanic category includes Hispanics of any race.

AIAN stands for American Indian or Alaska Native and NHOPI stands for Native Hawaiian or Other Pacific Islander.

Testing Coverage

Testing for the novel coronavirus is an important public health response to limiting the spread of the infection. Testing capacity was limited in Milwaukee County and across the country earlier in the epidemic, but then increased. Since the first case of COVID-19 was diagnosed in Milwaukee County on March 6, 2020, a total of 514130 COVID-19 tests have been performed, with 472559 negative results and 41571 positive results. This represents a positive test rate of 8.1% since the beginning of the epidemic.


As shown in **Figure 13**, the total number of tests performed per week increased until early July and then peaked and declined, with an increase observed in the last few weeks. As shown in **Figure 14**, the percentage of positive tests has varied over the course of the epidemic, with a high of 25-30% in early April. Since then, the percent positive has changed in tandem with expanded testing capacity. The percentage of positive tests was 11.1% over the past week compared to 10.4% the previous week. **Figure 14** also illustrates the 14-day trend in the percent positive tests, showing a significant increase.

Test result Number of tests performed 30000 Negative 25000 Positive 20000 15000 10000 5000 Lynos POLOS APTON bo bo bo ho ho ho ho ho ho One-week window start date


Figure 13: Milwaukee County number of tests per week

Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team

Figure 14: Milwaukee County percent positive tests



Data source: Wisconsin Electronic Disease Surveillance System (WEDSS) Created by the Milwaukee County COVID-19 Epidemiology Intel Team




Spatial Patterns of Cases and Testing


COVID-19 spread is spatially patterned. **Map 1** below illustrates the cumulative burden (all confirmed cases) of COVID-19 in Milwaukee County. **Map 2** shows cases confirmed over the last week. **Map 3** shows the overall testing rate across the population. **Map 4** shows the testing rate over the last week. **Map 5** depicts the percentage of tests that were confirmed positive. **Map 6** shows cumulative COVID-19 related hospitalizations. **Map 7** shows the percentage of cases who have been hospitalized. All are crude rate maps created using census block group level COVID-19 data from WEDSS and population data from the US Census. The maps are smoothed to protect confidentiality and ensure that rates are stable while still providing geographic detail. Deciles are used to define categories. High rates are depicted in red with lower rates depicted in blue. Of note, some of the higher rates observed can be attributed to infections that have spread within group quarters, such as a nursing home, prison, or long-term care facility.

Decile Map 1: All confirmed cases of COVID-19

Created by the Milwaukee County Covid-19 Epidemiology Intel Team




Decile Map 2: Confirmed cases of COVID-19 within the last week




Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 confirmed cases included.


Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Decile Map 3: Overall testing rate



Decile Map 4: Testing rate within the last week






Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 tests included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data) 2018 American Community Survey (population data) City of Milwaukee Map Milwaukee Portal (neighborhood boundaries) Census Bureau TIGER/Line Shapefiles (census place boundaries)

Created by the Milwaukee County Covid-19 Epidemiology Intel Team


Decile Map 5: Percentage of tests that were confirmed positive



Method: A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 positive tests included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)


Decile Map 6: COVID-19 related hospitalizations



Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 hospitalized cases included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Decile Map 7: Percentage of COVID-19 cases that were hospitalized



Method:
A grid of points is used to estimate rates continuously across the map, based on the nearest cases with a minimum of 15 confirmed cases included.

Data Sources: Wisconsin Electronic Disease Surveillance System (WEDSS) (incidence data)

Data Sources & Acknowledgments

This report was created by faculty and staff in the Medical College of Wisconsin (MCW) Institute for Health and Equity (IHE) in partnership with representatives from local health departments and faculty from the University of Wisconsin-Milwaukee Zilber School of Public Health. Data sources include the Wisconsin Electronic Disease Surveillance System (WEDSS), the US Census Bureau, the Milwaukee County Medical Examiner's office, the Emergency Medicine Resource, and publicly available data obtained from local health and emergency response agencies. Data from the Wisconsin Electronic Data Surveillance System (WEDSS) summarized for the week includes data from October 14, 2020 through October 20, 2020. This work was funded by the Advancing a Healthier Wisconsin Endowment at the Medical College of Wisconsin.

Contact Information

For additional questions on this report, please contact Darren Rausch, Health Officer/Director, Greenfield Health Department, and Lead, Milwaukee County COVID-19 Epidemiology Intel Team: Darren.Rausch@greenfieldwi.us or (414) 329-5275.